



Influence of two electroejaculation voltages on the seminal quality of alpacas (*Vicugna pacos*) Influencia de dos voltajes de electroeyaculación en la calidad seminal de alpacas (*Vicugna pacos*)

García-Díaz Juan Ramón^{1*}, Chicaiza Sánchez Luis Alonso², Garzón Jarrin Rafael Alfonso²,
Masabanda-Soria Eddy Bryan²

Article Data

¹Central University "Marta Abreu" of Las Villas.
Faculty of Agricultural Sciences.
Department of Veterinary Medicine and Animal
Husbandry.
Carretera a Camajuaní Km 5 ½ Santa Clara.
CP 54830, Santa Clara.
Villa Clara, Cuba.

²Technical University of Cotopaxi.
Faculty of Agricultural Sciences and Natural Re-
sources.
Av. Simón Rodríguez s/n Barrio El Ejido Sector San
Felipe.
Tel: (593) 03 2252205/2252307/2252346.
CAREN: 2266164
Latacunga - Ecuador.

*Contact address:

Central University "Marta Abreu" of Las Villas.
Faculty of Agricultural Sciences.
Department of Veterinary Medicine and Animal
Husbandry.
Carretera a Camajuaní Km 5 ½ Santa Clara.
CP 54830, Santa Clara.
Villa Clara, Cuba.

Juan Ramón García-Díaz

E-mail address: juanramon@uclv.edu.cu

Keywords:

Ejaculate,
volume,
motility,
sperm concentration,
sperm morphology,
alpacas.

J. Selva Andina Anim. Sci.
2022; 9(1):15-22.

Article ID: 109/JSAAS/2021

Article history.

Received November 2021.
Returned February 2022.
Accepted March 2022.
Available online, April 2022.

*Edited by: Selva Andina
Research Society*

Palabras clave:

Eyaculado,
volumen,
motilidad,
concentración espermática,
morfología espermática,
alpacas.

Abstract

The objective of the research was to evaluate the influence of two voltages in the electroejaculation protocols on the seminal quality of alpacas (*Vicugna pacos*). It was developed in the Reproductive Biotechnology Laboratory, Salache Academic Experimental Center, Technical University of Cotopaxi. Eight male alpacas were used and from each one two ejaculates were collected weekly, one using the electroejaculation protocol that used several 10 volt stimuli, and the other using a protocol where the stimuli were 12 volts. The volume of the ejaculated determined itself for direct observation in measuring glass, the concentration; sperm motility, viability and morphology by Neubauer chamber, DM4B YDM6B mass microscope at 10X, Eosin-Nigrosin staining and observation of semen-eosin staining, respectively. Volume, concentration of ejaculates, motility, seminal pathologies were compared according to the electroejaculation protocol by means of the Student's t test for independent samples. The volume of the ejaculates was 0.75 ± 0.20 and 1.10 ± 0.39 mL, the concentration 2.12 ± 1.93 and 2.25 ± 1.89 ($\times 10^6$), the sperm mass motility 81.25 ± 11.81 % and 80.00 ± 14.14 %, mortality 17.50 ± 9.57 and 15.00 ± 10.80 % and morphoanomalies 12.50 ± 6.45 and 13.50 ± 8.01 % in the animals that received electroejaculation with 10 and 12 volts, respectively. There were no significant differences ($p > 0.05$) in the seminal parameters evaluated according to the voltage used in the ejaculation protocol. It is concluded that the increase in voltage in the electroejaculation protocol had no influence on the seminal quality parameters of alpacas

2022. Journal of the Selva Andina Animal Science®. Bolivia. All rights reserved.

Resumen

La investigación tuvo como objetivo evaluar la influencia de dos voltajes en los protocolos de electroeyaculación en la calidad seminal de alpacas (*Vicugna pacos*). Se desarrolló en el laboratorio de Biotecnología de la Reproducción, Centro Experimental Académico Salache, Universidad Técnica de Cotopaxi. Se utilizaron 8 alpacas machos y de cada uno se colectaron dos eyaculados con una frecuencia semanal, uno mediante el protocolo EE que usó varios estímulos de 10 V, y en el otro mediante un protocolo donde los estímulos fueron de 12 V. El volumen del eyaculado se determinó por observación directa en probeta graduada, la concentración; motilidad, viabilidad y morfología espermática mediante cámara de Neubauer, microscopio masal DM4B Y DM6B a 10X, tinción Eosina-Nigrosina y observación de la tinción semen-eosina, respectivamente. Se compararon el volumen, la concentración de los eyaculados, motilidad, patologías seminales según el protocolo de EE mediante la prueba de t-Student para muestras independientes. El volumen de los eyaculados fue 0.75 ± 0.20 y 1.10 ± 0.39 mL, la concentración 2.12 ± 1.93 y 2.25 ± 1.89 ($\times 10^6$), la motilidad masal los espermatozoides 81.25 ± 11.81 % y 80.00 ± 14.14 %, la mortalidad 17.50 ± 9.57 y 15.00 ± 10.80 %, las morfoanomalías 12.50 ± 6.45 y 13.50 ± 8.01 % en los animales que recibieron la electroeyaculación con 10 y 12 V, respectivamente. No existieron diferencias significativas ($p > 0.05$) en los parámetros seminales evaluados según el voltaje utilizado en el protocolo de eyaculación. Se concluye que el incremento del voltaje en el protocolo de electroeyaculación no tuvo influencia en los parámetros de calidad seminal de alpacas.

2022. Journal of the Selva Andina Animal Science®. Bolivia. Todos los derechos reservados. 2022.

Introduction

Alpacas are domestic animals that belong to the New World group, the Camelidae family, they live mainly in South America^{1,2}, their breeding constitutes a source of income for the residents of Andean communities in Peru, Bolivia, Chile, Argentina and Ecuador³.

To accelerate its genetic improvement, it is necessary to use reproductive biotechnologies, which include, among others, collection, evaluation and conservation of semen⁴.

There are differences in seminal composition, and a lower concentration of spermatozoa in ejaculates collected by electroejaculation (EE) compared to the other method that involves artificial vagina⁵, and including sperm quality evaluation after defreezing which is lower in semen collected by EE⁶. However, it is a viable method to collect ejaculates in males that are not trained or wild species⁷.

EE is based on the application of electrical stimuli to the pelvic nerves through a probe inserted into the rectum, thus promoting protrusion of the penis, erection and finally ejaculation⁸.

EE for alpaca semen collection was used with very variable results, the ejaculates had little volume (0.1 to 0.5 mL) and high concentration⁹. Later in this species it was widely utilized EE¹⁰⁻¹², it is also used in vicuna¹³, llama¹⁴⁻¹⁶.

The voltage used in the EE influences the sperm quality, in alpacas most of the protocols used use 2 to 10 V, which avoids stress in the animals, to obtain good quality ejaculates, according to their volume, concentration and motility, provided that the males are handled correctly prior to the application^{10,12,17}.

In other species, different voltages are used in the EE, which influence the time to induce ejaculation and volume. In llamas, 2 to 10 V are used and

the ejaculate is obtained after 6 min^{14,18}, 3 to 10 V¹⁹ were also used with similar results.

In this species, in Bolivia, the ejaculation occurred with a voltage of 18 V, obtaining 0.9 mL of ejaculate²⁰, in Peru, 2 to 20 V were applied, penile erection occurred at 15 V, while ejaculation required 20 V, 1 mL of ejaculate was collected²¹. In the vicuna, between 2 and 12 V have been used, ejaculation occurred at 6 V¹³.

Information on the effect of the voltage used in EE on the seminal quality of alpacas is scarce, due to the lack of a reliable and reproducible methodology for semen collection. The purpose of this work was to evaluate the influence of two voltages in the EE protocol on the seminal quality of alpacas (*Vicugna pacos*).

Materials and methods

The research was carried out between May and August 2020, in the province of Cotopaxi, Lata cunga canton, located between 0° 59' 10" south latitude (LS) and 78° 37' 13" west longitude (LW), at 2739 meters above sea level, with an average temperature of 12 °C, relative humidity of 75 %, average annual rainfall between 500 and 1500 mm, clay soil²².

The facilities of the Salache Academic Experimental Center (CEASA), Reproductive Biotechnology Laboratory of the Veterinary Medicine, Technical University of Coto Paxi were used.

Eight male alpacas (*V. pacos*) were selected, aged between 3 and 5 years, with 64±2.2 kg of weight, without reproductive problems, of congenital defects, clinically healthy, with feeding based on natural pastures and water ad libitum. Two ejaculates were col-

lected from each of these animals with a weekly frequency with the same number of EE protocols for this purpose; they had the same characteristics.

The animals were supplied to the alpaca wool and genetic improvement project of the Technical University of Cotopaxi by the indigenous communities of the provinces of Cañar, Chimborazo, Cotopaxi and Imbabura, in Ecuador.

The ES (Figure 1) was performed according to the basic principles of the technique described by Director et al.¹⁸, to which the following adjustments were made. i) several 10 V stimuli were used, with electrical stimulation periods of 4 s, 1 s with rest intervals of 1 s, for 4-8 min ii) the stimuli were 12 V, with equal intervals and duration. An ElectroJac 5® electro-ejaculator (Ideal Instruments®, Lansing, USA) was used for this procedure.

Figure 1 Application of EE to alpacas in this research



The ejaculates were collected in 50 mL falcon tubes (Figure 2), protected by an external cover that maintained a temperature of 37 °C, and were transported to the CEASA Reproductive Biotechnology Laboratory in less than 1 h from collection. Samples contaminated with urine were not processed.

Ejaculate volume was determined by direct observation using a graduated cylinder. Seminal concentration, using the Neubauer chamber and motility using a DM4B and DM6B mass microscope (Leica Microsystems AG, USA), at 10X. Both parameters were determined according to the procedures described by Allauca et al.²³.

Sperm viability was determined by staining Eosin-Nigrosin, dead spermatozoa do not they take colorant

and appear pink, however, the live ones appear translucent²⁴.

Sperm morphology was evaluated by observing semen-eosin staining in a scanning electron microscope, the number of morphologically normal and abnormal sperm from according to the recommendations of Pérez et al.²⁵.

Statistical processing. Descriptive statistics were obtained for each variable. The volume, concentration of the ejaculates, motility and semen pathologies were compared according to the electroejaculation protocol using the t-Student test for independent samples. In all the processes, the statistical package Statgraphics Centurion²⁶ was used.

Figure 2 Collection of the ejaculate

Results

Table 1 shows the seminal variables evaluated of the EE with 10 and 12 V.

There were no significant differences ($p > 0.05$) in the seminal parameters evaluated according to the voltage used in the ejaculation protocol. However, the volume was greater numerically when the voltage was 12 V. In this group there was a tendency to decrease mortality and morphoanomalies (Table 2).

Table 1 Descriptive statistics of the seminal variables of the alpacas studied

Statisticians	Volume (mL)		Concentration ($\times 10^6$)		Mass motility (%)		Mortality (%)		Morphoanomalies (%)	
	10 V	12 V	10 V	12 V	10 V	12 V	12 V	12 V	12 V	12 V
□	0.75	1.10	2.12	2.25	81.25	80.00	17.50	15.00	12.50	13.75
SD	0.20	0.39	1.93	1.89	11.81	14.14	9.574	10.80	6.45	8.01
CV	27.75	35.59	90.87	84.13	14.54	17.67	54.71	72.00	51.63	58.29
Minimum	0.50	0.60	1.00	1.00	65.0	60.00	10.00	5.00	5.00	6.00
Maximum	1.00	1.50	5.00	5.00	90.00	90.00	30.00	30.00	20.00	25.00
Range	0.50	0.90	4.00	4.00	25.00	30.00	20.00	25.00	15.00	19.00

SD: Standard deviation. CV: Coefficient of variation

Table 2 Seminal parameters (\bar{X}) of alpacas with two voltages used in the electroejaculation protocol

Parameters	Groups		\pm EE	p-value
	10 V	12 V		
Volume (mL)	0.75	1.10	0.15	0.16
Concentration ($\times 10^6$)	2.12	2.25	0.95	0.92
Mass motility (%)	81.25	80.00	6.51	0.89
Mortality (%)	17.50	15.00	5.10	0.74
Morphoanomalies (%)	12.50	13.75	3.63	0.81

Discussion

The volume of ejaculates, sperm concentration, sperm mass motility, mortality and morphoanomalies are lower than those obtained in alpacas from Peru²⁷.

The volume was lower than that indicated by Trujillo Bravo²⁸, who obtained 1.48 to 1.91 in males of the Huacaya Breed from 3 to 5 years of age. This work confirms that alpacas have low volume in their ejaculates²⁹.

Although alpacas have low sperm concentration²⁹, the one obtained in this research indicates that the ejaculates have quality to carry out artificial insemination and from them more seminal doses can be obtained, a parameter that will depend on the dilution rate to be used²¹.

Individual motility in alpaca is not an indicator of the quality of semen, fresh or cryopreserved²⁷, however, according to the criteria of Allauca et al.²³, in this investigation the mass and individual motility describe

the ejaculates as good and confirm that South American camelids have little or no individual mobility²⁹.

Morphoanomalies are related to the size of the spermatozoa, affect fertilization capacity, and can be found in any segment of the spermatozoa^{30,31}. The percentage of spermatozoa with abnormal shapes did not differ according to the voltage used in the EE, which is motivated by the fact that these cells present great polymorphism and vary morphologically inter individually in each male of the South American camelids^{32,33}.

Mortality was lower than that obtained in a study, the collection of semen in alpacas, was done by vaginal aspiration and artificial vagina, the percentage of dead zoosperms was 24.7 and 29.2 %, respectively³⁴. Any EE protocol must maintain the viability of the largest number of possible spermatozoa, since this is positively related to acrosomal integrity in vivo and in vitro³⁵.

The causes of the differences in the seminal parameters determined in this work with the consultants may be motivated by the differences in the edaphoclimatic, physiological and production conditions⁸.

The volume, concentration, motility, sperm morphoanomalies vary according to the physiological state of the male, age, race, feeding, frequency collection and psychosexual separation³⁶.

It is concluded that the increase in the voltage level from 10 to 12 V in the EE protocol had no influence on the ejaculate and seminal quality parameters of alpacas (*V. pacos*)

Source of financing

We must emphasize that the financing of the research Influence of two voltages in the protocols of electroejaculation in the seminal quality of alpacas (*V. pacos*), was carried out with the researchers' own resources.

Conflicts of interest

The authors declare that there were no potential conflicts of interest with respect to the authorship and/or publication of this article.

Acknowledgments

We sincerely thank the Biotechnology Laboratory of the Technical University of Cotopaxi, Veterinary Medicine Career, especially the Director of the Salache Experimental Center (CEASA) and staff who work there.

Ethical considerations

The protocols used in the study adhere to the European standards indicated in https://ec.europa.eu/environment/chemicals/lab_animals/legislation_en.htm for the use of animals in research.

Authors' contribution to the article

Luis Alonso Chicaiza Sánchez, search for information, study design, sampling, writing of the draft. *Rafael Alfonso Garzón Jarrin*, information search, sample taking, writing and drafting of the document. *Eddy Bryan Masabanda Soria*, processing and analysis of samples in the laboratory. *Juan Ramón García Díaz*, writing and drafting of the document, revision, edition and revision of the manuscript.

Research limitations

It was not possible to establish and validate a methodology for obtaining semen of this species by electroejaculation because of the limited quantity of processed samples of them literature cited.

Literature cited

1. Quispe Peña E. Adaptaciones hematológicas de los camélidos sudamericanos que viven en zonas de elevadas altitudes. *Rev Complut Cienc Vet* 2011;5(1):1-26.
2. Avilés Esquivel DF, Montero-Recalde MA, Barros-Rodríguez M. Los camélidos sudamericanos: Productos y subproductos usados en la Región Andina. *Actas Iberoam Conserv Anim* 2018;11: 30-8.
3. Vilá B, Arzamendia Y. Camélidos sudamericanos: sus valores y aportes a las personas. *Sustain Sci* 2020. DOI: <https://doi.org/10.1007/s11625-020-00874-y>
4. Singh B, Mal G, Gautam SK, Mukesh M. Reproduction biotechnology in camelids. In: Singh B, Mal G, Gautam SK, Mukesh M, editors. *Advances in Animal Biotechnology*. Switzerland: Springer Nature Switzerland AG; 2019. p. 145-53. DOI: https://doi.org/10.1007/978-3-030-21309-1_13
5. Marco-Jiménez F, Vicente JS, Viudes-de-Castro MP. Seminal plasma composition from ejaculates collected by artificial vagina and electroejaculation in Guirra Ram. *Reprod Domest Anim* 2008; 43(4):403-8. DOI: <https://doi.org/10.1111/j.1439-0531.2007.00923.x>
6. Jiménez-Rabadán P, Ramón M, García-Álvarez O, Maroto-Morales A, del Olmo E, Pérez-Guzmán MD, et al. Effect of semen collection method (artificial vagina vs. electroejaculation), extender and centrifugation on post-thaw sperm quality of Blanca-Celtibérica buck ejaculates. *Anim Reprod Sci* 2012;132(1-2):88-95. DOI: <https://doi.org/10.1016/j.anireprosci.2012.04.005>
7. Santiago-Moreno J, Coloma MA, Dorado J, Pulido-Pastor A, Gómez-Guillamon F, Salas-Vega R, et al. Cryopreservation of Spanish ibex (*Capra pyrenaica*) sperm obtained by electroejaculation outside the rutting season. *Theriogenology* 2009; 71(8):1253-60. DOI: <https://doi.org/10.1016/j.theriogenology.2008.12.019>
8. Abraham MC, Verdier K, Båge R, Morrell JM. Semen collection methods in alpacas. *Vet Rec* 2017; 180(25):613-4. DOI: <https://doi.org/10.1136/vr.104074>
9. Rodriguez J, Huanca W, Ramos M, Vásquez M, Espinoza J. Biophysical and biochemical characteristics of alpaca semen after collection by electroejaculation. *Reprod Fertil Dev* 2012;25(1):272. DOI: <https://doi.org/10.1071/RDv25n1Ab249>
10. Ordoñez C, Ampuero E, Alarcón V, Franco E, Hanzen C, Cucho, H. Three extenders evaluation in cryopreservation of alpaca (*Vicugna pacos*) semen in pellets. *Spermova* 2015;5(1):119-23. DOI: <http://doi.org/10.18548/aspe/0002.27>
11. Choez K, Arriaga I, Terreros M, Condori R, Arroyo G, Huanca W. Characteristics of the alpaca semen obtained by electroejaculation and motility during cooling. *Spermova* 2015;5(1):42-6. DOI: <http://doi.org/10.18548/aspe/0002.9>
12. Masabanda Soria EB. Inseminación artificial de alpacas con semen colectado por electroeyaculador ajustado a 2 voltajes [tesis licenciatura]. [Latacunga]: Universidad Técnica de Cotopaxi; 2020 [citado 26 de mayo de 2021]. Recuperado a partir de: <http://repositorio.utc.edu.ec/handle/27000/6740>
13. Pacheco JI, Mamni RH, Deza HW. Collection of semen by electroejaculation held captive in vicuñas. *Spermova* 2011;1(1):131-3.
14. Giuliano S, Director A, Gambarotta M, Trasorras V, Miragaya M. Collection method, season and individual variation on seminal characteristics in the llama (*Lama glama*). *Anim Reprod Sci* 2008;104 (2-4):359-69. DOI: <https://doi.org/10.1016/j.anireprosci.2007.02.016>

15. Giuliano S, Carretero M, Gambarotta M, Neild D, Trasorras V, Pinto M, et al. Improvement of llama (*Lama glama*) seminal characteristics using collagenase. *Anim Reprod Sci* 2010;118(1):98-102. DOI: <https://doi.org/10.1016/j.anireprosci.2009.06.005>
16. Giuliano SM, Chaves MG, Trasorras VL, Gambarotta M, Neild D, Director A, et al. Development of an artificial insemination protocol in llamas using cooled semen. *Anim Reprod Sci* 2012;131(3-4):204-10. DOI: <https://doi.org/10.1016/j.anireprosci.2012.03.010>
17. Abril-Sánchez S, Freitas-de-Melo A, Giriboni J, Santiago-Moreno J, Ungerfeld R. Sperm collection by electroejaculation in small ruminants: A review on welfare problems and alternative techniques. *Anim Reprod Sci* 2019;205:1-9. DOI: <https://doi.org/10.1016/j.anireprosci.2019.03.023>
18. Director A, Giuliano S, Carretero M, Pinto M, Trasorras V, Miragaya M. Electroejaculation and seminal characteristics in llama (*Lama glama*). *J Camel Pract Res* 2007;14 (2):203-6.
19. Gallegos Ibarra ML. Determinación de las características macroscópicas, microscópicas y subpoblaciones espermáticas del semen de llama (*Lama glama*) colectado por electroeyaculación y vagina artificial [tesis licenciatura]. [Cusco]: Universidad Nacional de San Antonio Abad del Cusco; 2019 [citado 16 de septiembre de 2021]. Recuperado a partir de: <http://repositorio.concytec.gob.pe/handle/20.500.12390/1494>
20. Valle Zapata EM. Evaluación de dos técnicas de colección de semen en llamas (*Lama glama*) en la estación experimental de Choquenaira [tesis licenciatura]. [La Paz]: Universidad Mayor de San Andrés; 2013 [citado 16 de octubre de 2021]. Recuperado a partir de: <https://repositorio.umsa.bo/handle/123456789/4259?show=full>
21. Ccoiso Aguila R. Estimación de parámetros de motilidad y determinación de subpoblaciones espermáticas en semen de llama (*Lama glama*). por dos métodos de colección [tesis licenciatura]. [Cusco]: Universidad Nacional de San Antonio Abad del Cusco; 2019 [citado 6 de octubre de 2021]. Recuperado a partir de: <http://repositorio.unsaac.edu.pe/handle/20.500.12918/3790?locale-attribute=en>
22. Datos meteorológicos del parque nacional Coto-paxi, Ecuador [Internet]. Instituto nacional de meteorología e hidrografía. 2021 [citado 5 de octubre de 2021]. Recuperado a partir de: <https://www.inamhi.gob.ec>
23. Allauca P, Ugarelli A, Santiani A. Determinación del potencial de membrana mitocondrial mediante citometría de flujo durante el proceso de criopreservación de espermatozoides epididimarios de alpacas. *Rev Investig Vet Perú* 2019;30(1):288-98. DOI: <https://doi.org/10.15381/rivep.v30i1.15677>
24. Juárez J, Santiani A. Determinación del porcentaje de viabilidad espermática mediante citometría de flujo durante el proceso de criopreservación en espermatozoides obtenidos de epidídimo de alpaca. *Rev Investig Vet Perú* 2019;30(3):1175-83. DOI: <https://doi.org/10.15381/rivep.v30i3.16608>
25. Pérez MG, Zevallos J, Pérez UH. Recovery of alpaca sperm from vas deferens during the breeding season. *Spermova* 2014;4(2):139-44.
26. Centurion XVI Novedades y mejoras [Internet]. Statgraphics. 2018 [citado 5 de marzo de 2021]. Recuperado a partir de: <https://statgraphics.net/centurion-xvi-novedades-y-mejoras/>
27. Flores NH, Cucho H, Carretero MI, Ciprián R, Quispe H, Calderón N, et al. Dimethylformamide cryoprotectant effect on cryopreserved alpaca sperm motility (*Vicugna pacos*) evaluated by analysis system ISAS®. *Spermova* 2015;5(1):47-50. DOI: <https://doi.org/10.18548/aspe/0002.10>

28. Trujillo Bravo J. Estudio histológico del espermatozoide de alpacas y su correlación con las características microscópicas de calidad seminal en el fundo Ucrucancha-Cerro de Pasco 2019 [tesis licenciatura]. [Cerro de Pasco]: Universidad Nacional Daniel Alcides Carrión; 2019 [citado 6 de octubre de 2021]. Recuperado a partir de: <http://repositorio.undac.edu.pe/handle/undac/1490>
29. Miragaya M, Martínez Sarraague M, Casaretto M, Rubin de Celis E, Carretero I, Giuliano S. Assessment of apparent viscosity and analysis of rheological profiles in llama ejaculates. En: 16th International Congress on Animal Reproduction (ICAR); 2008; July 13-17; Budapest. Hungary.
30. Peña FJ, Saravia F, García-Herreros M, Núñez-Martínez I, Tapia JA, Johannisson A, et al. Identification of sperm morphometric subpopulations in two different portions of the boar ejaculate and its relation to postthaw quality. *J Androl* 2005;26(6):716-23. DOI: <https://doi.org/10.2164/jadrol.05030>
31. Gontarz A, Banaszewska D, Gryzińska M, Andrasz K. Differences in drone sperm morphometry and activity at the beginning and end of the season. *Turk J Vet Anim Sci* 2016;40(5):598-602. DOI: <https://doi.org/10.3906/vet-1511-6>
32. Casaretto C, Lombardo DM, Giuliano S, Gambarrutta M, Carretero MI, Miragaya MH. Morphometric analysis of llama (*Lama glama*) sperm head. *Andrologia* 2012;44(Suppl 1):424-30. DOI: <https://doi.org/10.1111/j.1439-0272.2011.01200.x>
33. Soler C, Sancho M, García A, Fuentes M, Núñez J, Cucho H. Ejaculate fractioning effect on llama sperm head morphometry as assessed by the ISAS(®) CASA system. *Reprod Domest Anim* 2014;49(1):71-8. DOI: <https://doi.org/10.1111/rda.12226>
34. Alarcón V, García W, Bravo W. Inseminación artificial de alpacas con semen colectado por aspiración vaginal y vagina artificial. *Rev Investig Vet Perú* 2012;23(1):58-64.
35. Pérez-Rosales M, Román B, Santiani A. Efecto de la criopreservación en la integridad acrosomal de espermatozoides viables de alpaca evaluada mediante citometría de flujo. *Rev Investig Vet Perú* 2020;31(3):e18172. DOI: <https://doi.org/10.15381/rivep.v31i3.18172>
36. Juyena NS, Vencato J, Pasini G, Vazzana I, Stelletta C. Alpaca semen quality in relation to different diets. *Reprod Fertil Dev* 2013;25(4):683-90. DOI: <https://doi.org/10.1071/RD12050>

Note from the Editor:

Journal of the Selva Andina Animal Science (JSAAS) remains neutral with respect to published jurisdictional claims maps and institutional affiliations.